



**Національний
університет
біоресурсів і
природокористування
України**

**Факультет
ветеринарної
медицини**

НДІ Здоров'я тварин



**«ЄДИНЕ ЗДОРОВ'Я – 2022»
Матеріали Міжнародної наукової конференції**



**22-24 вересня 2022 р.
НУБіП України, м. Київ**

UDC 591.69:598.617.2

**DIVERSITY OF PHEASANT PARASITIC AGENTS
(*PHASIANUS COLCHICUS L*) HELD IN CAPTIVITY IN THE CENTRAL
AREA OF THE REPUBLIC OF MOLDOVA**

Zamornea Maria, PhD in Biology

Rusu Ștefan, PhD in Biology

Erhan Dumitru, Doctor Habilitate in Biology, Research Professor

Chihai Oleg, PhD in Biology

Gliga Olesea, PhD in Biology

Botnaru Nicolai, PhD in Biology

Institute of Zoology, 1 Academiei Street, Chisinau, Republic of Moldova

Corresponding author: mariazamornea@gmail.com

The common pheasant (*Phasianus colchicus L.*) is the most important bird for the avian fauna of the Republic of Moldova, both in terms of numbers and distribution, as well as hunting perspectives.

The efficient and continuous exploitation of species for hunting purposes requires the most detailed knowledge of their way of life, the correlations between the

populations of this species, as well as their level of parasite infestation (Zamornea M. et al. 2017).

The investigations regarding the determination of the pheasant parasite species have been performed in the Parasitology and Helminthology Laboratory of the Institute of Zoology. Biological, samples were collected from the Central area of the Republic of Moldova in the period 2020-2022.

The following methods were used: the coproovoscopic methods (Fulleborn, Darling), the coprolarvoscopic methods (Popov, Baermann), partial parasitological investigations (after K. I. Skriabin) and successive washing were used. The collected material was further examined using the МБС-9 magnifier (ob.14x2) and the Novex Holland B ob microscope. 20-40 WF 10x Din / 20 mm.

In pheasants from the 127 samples collected, 14 parasite species were recorded: *Capillaria annulata* (Molin, 1858), *Syngamus tracheia* (Montagu, 1811), *Heterakis isolonche* (Linstow, 1906), *Ascaridia galli* (Schrank, 1788), *Heterakis gallinarum* (Schrank, 1788), *Trichostrongylus tenuis* (Mehlis, 1846), *Capillaria caudinflata* (Zeder, 1800), *Eimeria colchici* (Norton, 1967), *Eimeria duodenalis* (Norton, 1967), *Eimeria phasiani* (Tyzzer, 1929,) *Choanotaenia infundibulum* (Bloch, 1779), *Raillietina tetragona* (Molin, 1858), *Prosthogonimus ovatus* (Rud., 1803), *Raillietina echinobotrida* (Megnin, 1880), which were distributed in the 4 classes (*Trematoda*, *Cestoda*, *Nematoda*, *Conoidasida*), 7 families (*Prosthogonimidae*, *Davaineidae*, *Capillariidae*, *Syngamidae*, *Heterakidae*, *Trichostrongylidae*, *Eimeriidae*) and 8 genes (*Prosthogonimus*, *Raillietina*, *Capillaria*, *Syngamus*, *Heterakis*, *Ascaridia*, *Trichostrongylus*, *Eimeria*).

The research was carried out with the support of the institutional project - state program: DIVERSITY OF HEMATOPHAGOUS ARTHROPODS, ZOO- AND PHYTO-HELMINTHS, THEIR VULNERABILITY AND TOLERANCE STRATEGIES TO CLIMATIC FACTORS AND ELABORATION OF INNOVATIVE PROCEDURES FOR INTEGRATED CONTROL OF SPECIES WITH SOCIO-ECONOMIC VALUE: 20.80009.7007.12 F, 2020-2022.